

Synthesis of Heteroannulated Indolopyrazines through Domino N–H Palladium-Catalyzed/Metal-Free Oxidative C–H Bond Activation

Yuriy A. Kvashnin, Egor V. Verbitskiy,* Ekaterina F. Zhilina, Gennady L. Rusinov, Oleg N. Chupakhin, and Valery N. Charushin



Cite This: *ACS Omega* 2020, 5, 15681–15690



Read Online

ACCESS |




Metrics & More



Article Recommendations



Supporting Information



ABSTRACT: A convenient approach to [1,2,5]oxadiazolo[3',4':5,6]pyrazino[2,3-*b*]indoles and their heteroannulated analogues bearing various aryl substituents in the backbone has been developed. This synthetic protocol is based on Pd-catalyzed Buchwald–Hartwig and subsequent annulation by intramolecular oxidative cyclodehydrogenation. The photophysical properties for new polycycles have been measured.

INTRODUCTION

Indolo[2,3-*b*]pyrazines are significant structural motifs possessing many important applications such as in medicinal chemistry and material sciences (Figure 1). Namely, indolo[2,3-*b*]pyrazine derivatives display a wide range of promising

biological activities, such as antibacterial,¹ anticancer,² and antiviral activities.³ Many indolopyrazine derivatives have also been suggested as promising materials for organic electronics such as organic light-emitting diodes,⁴ solar cells,⁵ and organic transistor memory devices.⁶

There are five main routes for construction of the indolopyrazine core that are described in the literature, and they should be discussed to identify their weaknesses (Scheme 1). The most famous methods for the preparation of indolopyrazine consist in the condensation of isatin with *o*-phenylenediamine (route 1, Scheme 1)^{1,2c,6,7} or condensation of *in situ*-generated 2,3-diaminoindole with α -dicarbonyl compounds (route 2, Scheme 1).⁸ Generally, the main limitations of these methods are obtaining regioisomeric mixtures of indolopyrazines, when unsymmetrical *o*-phenylenediamines or α -dicarbonyl compounds are used. Buchwald–Hartwig amination followed by C–H activation from secondary amines and 2,3-dibromoquinoxaline in one pot (route 3, Scheme 1)⁹ and a two-step approach using Suzuki cross-coupling and subsequent annulation by Pd-catalyzed







Figure 1. Representative compound bearing the indolo[2,3-*b*]pyrazine core.

Received: April 28, 2020

Accepted: June 5, 2020

Published: June 18, 2020



Scheme 1. Synthetic Routes to the Formation of the Indolo[2,3-*b*]pyrazine ScaffoldScheme 2. Synthesis of 5*H*-[1,2,5]Oxadiazolo[3',4':5,6]pyrazino[2,3-*b*]indole (I) and its 5-Alkyl-Substituted Derivatives (II)Scheme 3. Domino Synthesis of 5-Phenyl-5*H*-[1,2,5]oxadiazolo[3',4':5,6]pyrazino[2,3-*b*]indole 3a

twofold C–N coupling reactions with primary amines (route 4, Scheme 1)¹⁰ afford indolo[2,3-*b*]pyrazines in high yield but they similarly do not possess regioselectivity. Furthermore, routes 3 and 4 have not corresponded the concept of the pot, atom, and step economic (PASE)¹¹ synthetic procedure because it requires using 2,3-di(chloro)bromopyrazines that is necessary for an additional introduction of halogens in the pyrazine ring. Another regioselective method for the synthesis of the indolopyrazines based on a sequential rhodium-catalyzed formal [3+3] cycloaddition and aromatization reaction of various diazoindolinimines with azirines.¹² This protocol requires a highly expensive and non-eco-friendly rhodium catalyst and infrequent raw materials. All these features eventually lead to the necessity of developing new PASE processes to synthesize indolopyrazine and their heteroanalogues that will be based on the metal-free oxidative C–H bond functionalization (so-called S_N^H reaction).¹³

It should be mentioned that [1,2,5]oxadiazolo[3,4-*b*]pyrazine (so-called “furanopyrazine”) exhibits higher electron-withdrawing character than similar nonannulated pyrazine analogues and can be readily involved into the metal-free oxidative C–H bond functionalization.¹⁴ Nowadays, there is only one example of the synthesis of unsubstituted 5*H*-

[1,2,5]oxadiazolo[3',4':5,6]pyrazino[2,3-*b*]indole (I) and its 5-alkyl-substituted derivatives (II) as a new family of effective inhibitors for the β -catenin/T-cell factor protein–protein interaction (Scheme 2).¹⁵

By the foregoing, furazanopyrazine derivatives have been selected as major precursors of novel aryl-substituted 5*H*-[1,2,5]oxadiazolo[3',4':5,6]pyrazino[2,3-*b*]indoles.

RESULTS AND DISCUSSION

Synthesis. It has been found that the reaction of 5-(2-bromophenyl)-[1,2,5]oxadiazolo[3,4-*b*]pyrazine **1** with aniline **2a** afforded 5-phenyl-5*H*-[1,2,5]oxadiazolo[3',4':5,6]pyrazino[2,3-*b*]indole **3a** in good yield (Scheme 3). The conditions of the domino reaction of **1** with aniline **2a** were optimized (Table 1). The ligand, palladium precursor, base, and solvent were varied. The results showed that tricyclohexylphosphine (PCy₃) often gave better yields than other ligands. In fact, up to 45% yield of **3a** was achieved by employment of PCy₃ as a ligand in combination with Pd(OAc)₂ as the Pd source (Table 1, entry 10).


With the optimized conditions in hand, we examined the scope of the twofold C–N annulation reaction of **1** with

Table 1. Optimization for the Synthesis of 3a

entry	Pd precursor (equiv)	ligand (20 mol %)	base (2.5 equiv)	solvent	yield 3a (%)
1	Pd(PPh ₃) ₄ (5 mol %)		K ₃ PO ₄	1,4-dioxane	0
2	Pd(PPh ₃) ₄ (10 mol %)		K ₃ PO ₄	1,4-dioxane	9
3	Pd(PPh ₃) ₂ Cl ₂ (10 mol %)	PPh ₃	K ₃ PO ₄	1,4-dioxane	15
4	Pd ₂ (dba) ₃ (10 mol %)	PCy ₃	K ₃ PO ₄	1,4-dioxane	28
5	Pd(OAc) ₂ (10 mol %)	XantPhos	K ₃ PO ₄	1,4-dioxane	35
6	Pd(OAc) ₂ (10 mol %)	XPhos	K ₃ PO ₄	1,4-dioxane	33
7	Pd(OAc) ₂ (10 mol %)	PCy ₃	DABCO	1,4-dioxane	0
8	Pd(OAc) ₂ (10 mol %)	PCy ₃	K ₂ CO ₃	1,4-dioxane	35
9	Pd(OAc) ₂ (10 mol %)	PCy ₃	K ₃ PO ₄	1,4-dioxane	40
10	Pd(OAc) ₂ (10 mol %)	PCy ₃	K ₃ PO ₄	toluene	45

various anilines **2b–i** and benzylamine **2j**. The results showed that the annulations gave the same yields for anilines bearing both electron-withdrawing and -donating substituents. However, the domino reactions with sterically crowded *ortho*-**2b,f** and *meta*-substituted **2c,g** anilines have resulted in the expected 5-aryl-5*H*-[1,2,5]oxadiazolo[3',4':5,6]pyrazino[2,3-*b*]indole **3** in lower yields than with unsubstituted **2a** and para-substituted **2d,h** anilines (Scheme 4). In contrast, the reactions of **1** with alkylamines, using our optimized conditions (Table 1, entry 10), resulted in the formation of side products which were difficult to separate from the main product. The appropriate result was only achieved for the synthesis of product **3j** derived from benzylamine in 59% yield. Structures of 5*H*-[1,2,5]oxadiazolo[3',4':5,6]pyrazino[2,3-*b*]indoles were verified by X-ray diffraction analysis performed for the single-crystal sample of 5-(*m*-tolyl)-substituted derivative **3i** (Figure 2), thus supporting the ¹H and ¹³C NMR spectroscopic data.

A feasible reaction mechanism is depicted in Scheme 3. The reaction between the **1** and aniline **2a** proceeds through the Buchwald–Hartwig amination, which gives intermediate **A**, followed by an intramolecular nucleophilic attack at C(6)


Figure 2. ORTEP of **3i** with thermal ellipsoids at the 50% probability level.



position furazanopyrazine core **B**. The subsequent oxidation of dihydropyrazine intermediate **B** by air oxygen leads to the polycyclic system **3a**.

To confirm this hypothesis, we carried out the reactions of furazanopyrazine **1** with aniline **2a** without using any palladium catalyst only in the presence of various bases. It has been shown that transition-metal-free oxidative dehydrogenation cross-coupling reaction leads to **4a** with the highest yield of 55% in the presence of sodium hydride as a base (Scheme 5). Using these conditions allowed us to obtain a wide range of corresponding S_N^H products by the reaction of **1** with the same series of amines. It is supposed that these reactions' C–H functionalization proceeds through nucleophilic substitution of hydrogen according to the two-step “Addition–Oxidation” mechanism.¹³


To increase the yields of polycyclic compounds **3**, we have carried out the annulation reaction of **4a** in the earlier developed best conditions of Buchwald–Hartwig amination (Scheme 6). Unfortunately, despite yields of intramolecular C–N coupling annulation being 72%, the two-step overall yield of **3a** from compound **1** to **3a** achieved only to 40%.

The suggested approach to the construction of the indolo[2,3-*b*]pyrazine core has also been applied successfully for the synthesis of different indolopyrazines with annulated benzene and/or thiophene rings (Scheme 7). In this regard, it is noteworthy that the only reaction of 5-(3-bromothiophen-2-yl)-[1,2,5]oxadiazolo[3,4-*b*]pyrazine **5a** with aniline **2a** yielded

Scheme 4. Synthesis and Structures of 5*H*-[1,2,5]Oxadiazolo[3',4':5,6]pyrazino[2,3-*b*]indole Derivatives **3a–j**

Scheme 5. Synthesis and Structures of *N*-Aryl-6-(2-bromophenyl)-[1,2,5]oxadiazolo[3,4-*b*]pyrazin-5-amines 4a–iScheme 6. Synthesis of 5-Phenyl-5*H*-[1,2,5]oxadiazolo[3',4':5,6]pyrazino[2,3-*b*]indole 3a by Intramolecular Buchwald–Hartwig Amination

Scheme 7. Synthesis of Heteroannulated Indolopyrazine Derivatives 7a–d



desired 8-phenyl-8*H*-[1,2,5]oxadiazolo[3,4-*b*]thieno-[2',3':4,5]pyrrolo[2,3-*e*]pyrazine 7a and compound 6a as a by-product. In other cases, the reactions of the bromine-

substituted derivatives 5b–d with aniline 2a led to the classic Buchwald–Hartwig amination products 6b–d in 77–89% yields. Notably, using PCy₃ as a ligand for the palladium

catalyst in this case afforded products in yields only up to 40%, while the replacement of it by Xantphos significantly increased the yield of compounds **6b–d**.

Further treatment of amination products **6b–d** by acids, followed by oxidation with air oxygen, also leads to similar cyclization products **7b–d**. Following the literature, trifluoroacetic acid¹⁶ was used for the cyclization of thiophenyl-substituted quinoxaline **6b**, while a mixture of hydrochloric acid¹⁷ in ethanol led to the transformation of amino derivatives **6a,c** into indolopyrazines **7c,b** in high yields (Scheme 7).

CONCLUSIONS

In summary, we have described a convenient approach to construct derivatives of a key heterocyclic system, namely, [1,2,5]oxadiazolo[3',4':5,6]pyrazino[2,3-*b*]indoles, bearing various substituents in the pyrrole fragment, based on domino Pd-catalyzed Buchwald–Hartwig and subsequent annulation by intramolecular oxidative cyclodehydrogenation reactions. The two-step synthesis of other heteroannulated indolopyrazine analogues has been realized using this synthetic approach. The further design and synthesis of different indoloannulated derivatives using this reaction procedure, as well as the elucidation of photophysical and electronic properties of these polycyclic compounds, are currently in progress.

EXPERIMENTAL SECTION

General Information. All reagents and solvents were obtained from commercial sources and dried using standard procedures before use. ¹H and ¹³C NMR spectra were obtained on Bruker DRX-400, AVANCE-500, and AVANCE-600 spectrometers with TMS as internal standards. Elemental analysis was carried out on a Eurovector EA 3000 automated analyzer. Melting points were determined on Boetius combined heating stages and were not corrected. All solvents used were dried and distilled as per standard procedures. IR spectra of samples (solid powders) were recorded on a Spectrum One Fourier transform IR spectrometer (PerkinElmer) equipped with a diffuse reflectance attachment (DRA). UV–vis spectra were recorded for ~10^{−5} to 10^{−6} M CH₂Cl₂ solutions with a Shimadzu UV-2401PC spectrophotometer. X-ray diffraction analysis was performed on an automated X-ray diffractometer “Xcalibur E” based on the standard procedure. The deposition number CCDC 1989238 for **3i** contains the Supporting Information crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

General Procedure for the Synthesis of 2-Bromophenyl and 3-(Bromothiophen-2-yl)-Substituted 1,4-Diazine Derivatives (1 and 5a–d). Selenium dioxide (1.1 g, 10 mmol) was dissolved in 1,4-dioxane/H₂O 15:1 (16 mL) and the mixture was heated at reflux for 5 min. 2'-Bromoacetophenone [or 2-acetyl-3-bromothiophene] (10 mmol) was added and heating continued for next 12 h. The reaction mixture was filtered and the filtrate was concentrated under reduced pressure to give the corresponding aryl glyoxal as light-yellow oil. The solution of 3,4-diaminofurazan [or 1,2-phenylenediamine] (10 mmol) and crude aryl glyoxal in a mixture of EtOH (5 mL) and CH₃COOH (5 mL) was refluxed for 1 h. After that, the mixture was cooled down, and

the precipitate was filtered off and washed with ethanol and then air-dried.

5-(2-Bromophenyl)-[1,2,5]oxadiazolo[3,4-*b*]pyrazine (1). 2'-Bromoacetophenone and 3,4-diaminofurazan were used for the synthesis of product **1**.

Yellow powder; yield 2.19 g (79%); mp 135 °C. ¹H NMR (500 MHz, DMSO-*d*₆): δ 9.44 (s, 1H), 7.91 (dd, *J* = 8.0, 1.1 Hz, 1H), 7.79 (dd, *J* = 7.6, 1.7 Hz, 1H), 7.68 (td, *J* = 7.6, 1.1 Hz, 1H), 7.63–7.59 (m, 1H). ¹³C NMR (126 MHz, DMSO-*d*₆): δ 162.4, 157.1, 151.9, 151.1, 136.1, 133.4, 133.1, 132.4, 128.5, 121.0. Anal. Calcd for C₁₀H₅BrN₄O (277.08): C, 43.35; H, 1.87; N, 20.22. Found: C, 43.41; H, 1.84; N, 20.10.

5-(3-Bromothiophen-2-yl)-[1,2,5]oxadiazolo[3,4-*b*]pyrazine (5a). 2-Acetyl-3-bromothiophene and 3,4-diaminofurazan were used for the synthesis of product **5a**.

Pale yellow powder; yield 2.15 g (76%); mp 174–176 °C. ¹H NMR (500 MHz, DMSO-*d*₆): δ 9.79 (s, 1H), 8.19 (d, *J* = 5.3 Hz, 1H), 7.46 (d, *J* = 5.3 Hz, 1H). ¹³C NMR (126 MHz, DMSO-*d*₆): δ 154.5, 154.0, 151.9, 151.1, 135.3, 134.2, 134.1, 114.8. Anal. Calcd for C₈H₃BrN₄OS (283.10): C, 33.94; H, 1.07; N, 19.79. Found: C, 33.92; H, 0.96; Br, 28.22; N, 19.69.

2-(3-Bromothiophen-2-yl)quinoxaline (5b). 2-Acetyl-3-bromothiophene and 1,2-phenylenediamine were used for the synthesis of product **5b**.

Beige powder; yield 2.39 g (82%); mp 111–112 °C. ¹H NMR (500 MHz, DMSO-*d*₆): δ 9.72 (s, 1H), 8.15–8.09 (m, 2H), 7.98 (d, *J* = 5.3 Hz, 1H), 7.93–7.88 (m, 2H), 7.36 (d, *J* = 5.3 Hz, 1H). ¹³C NMR (126 MHz, DMSO-*d*₆): δ 146.3, 142.7, 141.3, 140.7, 135.8, 132.8, 131.3, 131.1, 130.4, 128.9, 128.8, 109.8. Anal. Calcd for C₁₂H₇BrN₂S (291.17): C, 49.50; H, 2.42; N, 9.62. Found: C, 49.54; H, 2.56; N, 9.38.

2-(2-Bromophenyl)pyrazine (5c). Commercially available material **5c** was used in our experiments.

2-(2-Bromophenyl)quinoxaline (5d). 2'-Bromoacetophenone and 1,2-phenylenediamine were used for the synthesis of product **5b**.

Pale yellow powder; yield 2.28 g (77%); mp 112–113 °C (ref 18 116–118 °C). ¹H NMR (400 MHz, CDCl₃): δ 9.20 (s, 1H), 8.21 (td, *J* = 6.1, 3.2 Hz, 2H), 7.85 (dt, *J* = 6.4, 3.4 Hz, 2H), 7.76 (dd, *J* = 8.0, 1.2 Hz, 1H), 7.69 (dd, *J* = 7.7, 1.8 Hz, 1H), 7.52 (td, *J* = 7.5, 1.2 Hz, 1H), 7.39 (td, *J* = 7.7, 1.7 Hz, 1H). Anal. Calcd for C₁₄H₉BrN₂ (285.14): C, 58.97; H, 3.18; N, 9.82. Found: C, 58.88; H, 3.38; N, 9.69. Data similar to the literature.¹⁸

General Procedure for the Synthesis of 5-Aryl-5H-[1,2,5]oxadiazolo[3',4':5,6]pyrazino[2,3-*b*]indoles (3a–j). A stirred mixture of 5-(2-bromophenyl)-[1,2,5]oxadiazolo[3,4-*b*]pyrazine (**1**) (135 mg, 0.5 mmol), corresponding anilines (0.6 mmol), tricyclohexylphosphine (28 mg, 20 mol %), Pd(OAc)₂ (11 mg, 10 mol %), and K₃PO₄ (265 mg, 1.25 mmol) in degassed 1,4-dioxane (15 mL) was heated at reflux under nitrogen for 15 h in a Schlenk tube. The reaction mixture was cooled, filtered, acidified with concentrated CH₃COOH (1 mL), and dissolved with a mixture of EtOAc and water 1:1 (50 mL), and the organic layer was separated. The aqueous layer was extracted with EtOAc (2 × 25 mL). The combined organic extracts were dried with MgSO₄ and the solvents were evaporated. The organic layer was separated, dried over anhydrous Na₂SO₄, and concentrated. The crude product was purified by column chromatography using hexane and ethyl acetate as an eluent.

5-Phenyl-5H-[1,2,5]oxadiazolo[3',4':5,6]pyrazino[2,3-*b*]indole (3a). Red powder; yield 65 mg (45%); mp 285–286

$^{\circ}\text{C}$. R_f 0.45 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 8.41–8.38 (m, 1H), 7.83 (ddd, $J = 8.4$, 7.5, 1.3 Hz, 1H), 7.72–7.70 (m, 4H), 7.62 (ddt, $J = 6.5$, 4.9, 3.3 Hz, 1H), 7.50 (td, $J = 7.6$, 0.8 Hz, 1H), 7.32 (d, $J = 8.2$ Hz, 1H). ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$): δ 153.2, 151.7, 151.5, 148.8, 135.8, 133.5, 130.0, 129.0, 127.1, 124.8, 123.6, 118.3, 111.4. IR (DRA): 434, 446, 466, 496, 565, 592, 617, 627, 666, 691, 749, 775, 796, 807, 857, 880, 915, 946, 1008, 1019, 1030, 1057, 1081, 1110, 1168, 1180, 1128, 1259, 1307, 1330, 1379, 1445, 1486, 1502, 1551, 1586, 1598, 1621, 3079, 3099 cm^{-1} . Anal. Calcd for $\text{C}_{16}\text{H}_9\text{N}_5\text{O}$ (287.28): C, 66.89; H, 3.16; N, 24.38. Found: C, 66.81; H, 3.18; N, 24.35.

5-(2-Methoxyphenyl)-5H-[1,2,5]oxadiazolo[3',4':5,6]-pyrazino[2,3-b]indole (3b). Red powder; yield 22 mg (14%); mp 248–250 $^{\circ}\text{C}$. R_f 0.31 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ 8.38 (dd, $J = 7.7$, 1.2 Hz, 1H), 7.80 (ddd, $J = 8.4$, 7.4, 1.3 Hz, 1H), 7.67–7.63 (m, 1H), 7.61 (dd, $J = 7.7$, 1.7 Hz, 1H), 7.48 (td, $J = 7.5$, 0.9 Hz, 1H), 7.41 (dd, $J = 8.5$, 1.2 Hz, 1H), 7.25 (td, $J = 7.6$, 1.2 Hz, 1H), 7.05 (d, $J = 8.1$ Hz, 1H), 3.77 (s, 3H). ^{13}C NMR (151 MHz, $\text{DMSO}-d_6$): δ 155.7, 153.3, 152.2, 152.1, 149.5, 136.4, 131.8, 130.2, 125.1, 123.9, 121.7, 118.5, 113.7, 112.5, 56.3. Anal. Calcd for $\text{C}_{17}\text{H}_{11}\text{N}_5\text{O}_2$ (317.30): C, 64.35; H, 3.49; N, 22.07. Found: C, 64.42; H, 3.45; N, 21.93.

5-(3-Methoxyphenyl)-5H-[1,2,5]oxadiazolo[3',4':5,6]-pyrazino[2,3-b]indole (3c). Red powder; yield 76 mg (48%); mp 258–260 $^{\circ}\text{C}$. R_f 0.25 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ 8.42–8.37 (m, 1H), 7.83 (ddd, $J = 8.4$, 7.4, 1.3 Hz, 1H), 7.63 (t, $J = 8.3$ Hz, 1H), 7.50 (t, $J = 7.5$ Hz, 1H), 7.36 (d, $J = 8.2$ Hz, 1H), 7.28 (dt, $J = 3.6$, 1.4 Hz, 2H), 7.21 (ddd, $J = 8.4$, 2.5, 1.0 Hz, 1H), 3.85 (s, 3H). ^{13}C NMR (151 MHz, $\text{DMSO}-d_6$): δ 153.7, 152.2, 152.1, 152.0, 149.3, 136.4, 135.0, 131.3, 125.3, 124.1, 119.7, 118.8, 115.2, 113.4, 112.1, 56.0. Anal. Calcd for $\text{C}_{17}\text{H}_{11}\text{N}_5\text{O}_2$ (317.30): C, 64.35; H, 3.49; N, 22.07. Found: C, 64.30; H, 3.58; N, 22.09.

5-(4-Methoxyphenyl)-5H-[1,2,5]oxadiazolo[3',4':5,6]-pyrazino[2,3-b]indole (3d). Orange powder; yield 95 mg (60%); mp 271–273 $^{\circ}\text{C}$. R_f 0.18 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (500 MHz, $\text{DMSO}-d_6$): δ 8.37 (ddd, $J = 7.7$, 1.3, 0.6 Hz, 1H), 7.81 (ddd, $J = 8.5$, 7.4, 1.3 Hz, 1H), 7.63–7.59 (m, 2H), 7.48 (td, $J = 7.6$, 0.8 Hz, 1H), 7.27–7.23 (m, 3H), 3.89 (s, 3H). ^{13}C NMR (126 MHz, $\text{DMSO}-d_6$): δ 159.4, 153.2, 151.7, 151.5, 149.2, 135.8, 128.6, 125.9, 124.7, 123.4, 118.2, 115.2, 111.4, 55.5. IR (DRA): 453, 464, 484, 529, 567, 592, 623, 635, 644, 694, 725, 755, 771, 788, 797, 804, 815, 835, 842, 852, 875, 911, 936, 953, 969, 987, 1007, 1015, 1028, 1058, 1111, 1155, 1180, 1232, 1263, 1304, 1326, 1384, 1445, 1465, 1488, 1518, 1552, 1595, 1610, 1619, 3009, 3062 cm^{-1} . Anal. Calcd for $\text{C}_{17}\text{H}_{11}\text{N}_5\text{O}_2$ (317.30): C, 64.35; H, 3.49; N, 22.07. Found: C, 64.51; H, 3.54; N, 22.26.

5-(3,4-Dimethoxyphenyl)-5H-[1,2,5]oxadiazolo[3',4':5,6]-pyrazino[2,3-b]indole (3e). Red powder; yield 111 mg (64%); mp 271–272 $^{\circ}\text{C}$. R_f 0.28 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (500 MHz, $\text{DMSO}-d_6$): δ 8.37 (dt, $J = 7.7$, 0.9 Hz, 1H), 7.81 (ddd, $J = 8.4$, 7.4, 1.3 Hz, 1H), 7.48 (td, $J = 7.5$, 0.8 Hz, 1H), 7.30 (d, $J = 8.1$ Hz, 1H), 7.26 (dd, $J = 5.4$, 3.2 Hz, 2H), 7.22 (dd, $J = 8.5$, 2.2 Hz, 1H), 3.89 (s, 3H), 3.78 (s, 3H). ^{13}C NMR (126 MHz, $\text{DMSO}-d_6$): δ 153.2, 151.8, 151.5, 149.5, 149.4, 149.2, 135.8, 125.9, 124.6, 123.4, 119.8, 118.1, 112.3, 111.6, 110.9, 55.8. Anal. Calcd for $\text{C}_{18}\text{H}_{13}\text{N}_5\text{O}_3$ (347.33): C, 62.24; H, 3.77; N, 20.16. Found: C, 62.12; H, 3.61; N, 20.08.

5-(2-Nitrophenyl)-5H-[1,2,5]oxadiazolo[3',4':5,6]-pyrazino[2,3-b]indole (3f). Orange powder; yield 17 mg

(10%); mp 274–275 $^{\circ}\text{C}$. R_f 0.45 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ 8.51–8.36 (m, 2H), 8.12 (td, $J = 7.7$, 1.5 Hz, 1H), 7.98 (ddd, $J = 8.3$, 6.5, 1.6 Hz, 2H), 7.89–7.83 (m, 1H), 7.55 (t, $J = 7.5$ Hz, 1H), 7.29 (d, $J = 8.2$ Hz, 1H). ^{13}C NMR (151 MHz, $\text{DMSO}-d_6$): δ 153.1, 152.1, 151.9, 151.8, 148.8, 146.4, 136.7, 136.4, 132.0, 131.6, 127.0, 126.9, 125.5, 124.7, 118.8, 112.1. Anal. Calcd for $\text{C}_{16}\text{H}_8\text{N}_6\text{O}_3$ (332.27): C, 57.84; H, 2.43; N, 25.29. Found: C, 57.89; H, 2.34; N, 25.17.

5-(3-Nitrophenyl)-5H-[1,2,5]oxadiazolo[3',4':5,6]-pyrazino[2,3-b]indole (3g). Orange powder; yield 56 mg (34%); mp 287–290 $^{\circ}\text{C}$ (decomp.). R_f 0.36 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ 8.60 (t, $J = 2.2$ Hz, 1H), 8.48 (ddd, $J = 8.3$, 2.3, 1.0 Hz, 1H), 8.43 (dd, $J = 7.7$, 1.2 Hz, 1H), 8.23 (ddd, $J = 7.9$, 2.1, 1.0 Hz, 1H), 8.03 (t, $J = 8.1$ Hz, 1H), 7.86 (ddd, $J = 8.5$, 7.4, 1.3 Hz, 1H), 7.54 (td, $J = 7.5$, 0.8 Hz, 1H), 7.47 (d, $J = 8.2$ Hz, 1H). ^{13}C NMR (151 MHz, $\text{DMSO}-d_6$): δ 153.7, 152.2, 152.1, 152.0, 149.1, 148.7, 136.4, 135.0, 134.3, 132.1, 125.4, 124.5, 124.3, 122.9, 119.0, 112.0. Anal. Calcd for $\text{C}_{16}\text{H}_8\text{N}_6\text{O}_3$ (332.27): C, 57.84; H, 2.43; N, 25.29. Found: C, 57.80; H, 2.25; N, 25.18.

5-(4-Nitrophenyl)-5H-[1,2,5]oxadiazolo[3',4':5,6]-pyrazino[2,3-b]indole (3h). Orange powder; yield 91 mg (55%); mp 271–273 $^{\circ}\text{C}$. R_f 0.35 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ 8.60–8.56 (m, 2H), 8.44 (dd, $J = 7.8$, 1.2 Hz, 1H), 8.09–8.05 (m, 2H), 7.88 (ddd, $J = 8.5$, 7.4, 1.3 Hz, 1H), 7.57–7.52 (m, 2H). ^{13}C NMR (151 MHz, $\text{DMSO}-d_6$): δ 153.8, 152.2, 152.1, 152.0, 148.2, 147.2, 139.7, 136.4, 128.5, 125.8, 125.5, 124.7, 119.2, 112.2. IR (DRA): 463, 477, 492, 529, 598, 623, 689, 709, 747, 770, 793, 804, 827, 853, 865, 885, 913, 944, 1012, 1025, 1044, 1061, 1109, 1159, 1174, 1228, 1296, 1324, 1353, 1377, 1421, 1446, 1464, 1487, 1504, 1524, 1552, 1590, 1614, 3077, 3118 cm^{-1} . Anal. Calcd for $\text{C}_{16}\text{H}_8\text{N}_6\text{O}_3$ (332.27): C, 57.84; H, 2.43; N, 25.29. Found: C, 57.54; H, 2.42; N, 25.15.

5-(*m*-Tolyl)-5H-[1,2,5]oxadiazolo[3',4':5,6]pyrazino[2,3-b]indole (3i). Red powder; yield 93 mg (62%); mp 262–264 $^{\circ}\text{C}$. R_f 0.47 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 8.38 (d, $J = 7.6$ Hz, 1H), 7.87–7.77 (m, 1H), 7.60 (t, $J = 7.7$ Hz, 1H), 7.52–7.46 (m, 3H), 7.43 (d, $J = 7.6$ Hz, 1H), 7.31 (d, $J = 8.2$ Hz, 1H), 2.46 (s, 3H). ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$): δ 153.2, 151.7, 151.5, 148.8, 139.6, 135.8, 133.4, 129.8, 129.7, 127.4, 124.7, 124.2, 123.5, 118.3, 111.5, 20.9. Anal. Calcd for $\text{C}_{17}\text{H}_{11}\text{N}_5\text{O}$ (301.30): C, 67.77; H, 3.68; N, 23.24. Found: C, 67.88; H, 3.65; N, 23.41.

5-Benzyl-5H-[1,2,5]oxadiazolo[3',4':5,6]pyrazino[2,3-b]indole (3j). Red powder; yield 89 mg (59%); mp 210–212 $^{\circ}\text{C}$. R_f 0.17 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ 8.35–8.30 (m, 1H), 7.81 (td, $J = 7.8$, 7.4, 1.3 Hz, 1H), 7.57 (d, $J = 8.2$ Hz, 1H), 7.50–7.44 (m, 2H), 7.42 (t, $J = 7.5$ Hz, 1H), 7.34 (dd, $J = 8.2$, 6.6 Hz, 2H), 7.31–7.27 (m, 1H), 5.55 (s, 2H). ^{13}C NMR (151 MHz, $\text{DMSO}-d_6$): δ 153.6, 152.3, 152.3, 151.9, 148.5, 136.1, 136.0, 129.2, 128.2, 128.0, 125.2, 123.6, 118.8, 112.2, 44.9. Anal. Calcd for $\text{C}_{17}\text{H}_{11}\text{N}_5\text{O}$ (301.30): C, 67.77; H, 3.68; N, 23.24. Found: C, 67.56; H, 3.81; N, 23.24.

General Procedure for the Synthesis of *N*-Aryl-6-(2-bromophenyl)-[1,2,5]oxadiazolo[3,4-*b*]pyrazin-5-amines (4a–i). A stirred mixture of 5-(2-bromophenyl)-[1,2,5]oxadiazolo[3,4-*b*]pyrazine (1) (135 mg, 0.5 mmol), corresponding anilines (0.6 mmol), and sodium hydride (24 mg, 1.0 mmol) in dry 1,4-dioxane (5 mL) was stirred at room temperature for 48 h. The reaction mixture was acidified with

concentrated CH_3COOH (1 mL). The solvent was distilled off *in vacuo*, and the crude product was purified by column chromatography using hexane and ethyl acetate as an eluent.

6-(2-Bromophenyl)-N-phenyl-[1,2,5]oxadiazolo[3,4-*b*]pyrazin-5-amine (4a). Yellow powder; yield 65 mg (55%); mp 174–175 °C. R_f 0.26 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ 9.43 (s, 1H), 7.85 (dd, $J = 8.1$, 1.1 Hz, 1H), 7.76 (dd, $J = 7.6$, 1.7 Hz, 1H), 7.67 (td, $J = 7.6$, 1.2 Hz, 1H), 7.63–7.57 (m, 3H), 7.46–7.40 (m, 2H), 7.26 (tt, $J = 7.3$, 1.2 Hz, 1H). ^{13}C NMR (151 MHz, $\text{DMSO}-d_6$): δ 161.4, 152.4, 152.3, 150.2, 137.8, 133.5, 132.9, 132.0, 130.5, 129.0, 128.9, 127.6, 126.4, 124.8, 121.7. Anal. Calcd for $\text{C}_{16}\text{H}_{10}\text{BrN}_5\text{O}$ (368.19): C, 52.19; H, 2.74; N, 19.02. Found: C, 52.37; H, 2.63; N, 18.97.

6-(2-Bromophenyl)-N-(2-methoxyphenyl)-[1,2,5]-oxadiazolo[3,4-*b*]pyrazin-5-amine (4b). Orange powder; yield 108 mg (54%); mp 207–208 °C. R_f 0.42 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ 8.43 (s, 1H), 8.35–8.28 (m, 1H), 7.96 (d, $J = 8.1$ Hz, 1H), 7.82 (dd, $J = 7.5$, 1.7 Hz, 1H), 7.75 (t, $J = 7.5$ Hz, 1H), 7.68 (td, $J = 7.8$, 1.7 Hz, 1H), 7.27–7.20 (m, 1H), 7.14–7.06 (m, 2H), 3.68 (s, 3H). ^{13}C NMR (151 MHz, $\text{DMSO}-d_6$): δ 161.2, 152.4, 151.2, 150.4, 150.1, 134.8, 133.7, 133.4, 131.6, 129.3, 126.8, 126.7, 122.0, 121.5, 121.3, 111.9, 56.7. Anal. Calcd for $\text{C}_{17}\text{H}_{12}\text{BrN}_5\text{O}_2$ (398.21): C, 51.27; H, 3.04; N, 17.59. Found: C, 51.50; H, 3.12; N, 17.81.

6-(2-Bromophenyl)-N-(3-methoxyphenyl)-[1,2,5]-oxadiazolo[3,4-*b*]pyrazin-5-amine (4c). Yellow powder; yield 115 mg (58%); mp 171–172 °C. R_f 0.32 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ 9.36 (s, 1H), 7.85 (dd, $J = 8.1$, 1.1 Hz, 1H), 7.75 (dd, $J = 7.6$, 1.7 Hz, 1H), 7.66 (td, $J = 7.6$, 1.2 Hz, 1H), 7.59 (ddd, $J = 8.2$, 7.4, 1.7 Hz, 1H), 7.34 (t, $J = 8.1$ Hz, 1H), 7.27–7.19 (m, 2H), 6.85 (ddd, $J = 8.3$, 2.6, 1.0 Hz, 1H), 3.76 (s, 3H). ^{13}C NMR (151 MHz, $\text{DMSO}-d_6$): δ 161.3, 159.7, 152.3, 152.2, 150.2, 139.0, 135.8, 133.5, 132.9, 132.1, 129.8, 128.9, 121.8, 117.0, 111.6, 110.6, 55.7. Anal. Calcd for $\text{C}_{17}\text{H}_{12}\text{BrN}_5\text{O}_2$ (398.21): C, 51.27; H, 3.04; N, 17.59. Found: C, 51.40; H, 2.97; N, 17.32.

6-(2-Bromophenyl)-N-(4-methoxyphenyl)-[1,2,5]-oxadiazolo[3,4-*b*]pyrazin-5-amine (4d). Red powder; yield 125 mg (63%); mp 117–119 °C. R_f 0.35 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (500 MHz, $\text{DMSO}-d_6$): δ 9.34 (s, 1H), 7.84 (dd, $J = 8.0$, 1.1 Hz, 1H), 7.74 (dd, $J = 7.6$, 1.7 Hz, 1H), 7.65 (td, $J = 7.5$, 1.1 Hz, 1H), 7.58 (td, $J = 7.8$, 1.8 Hz, 1H), 7.53–7.45 (m, 2H), 7.02–6.97 (m, 2H), 3.77 (s, 3H). ^{13}C NMR (151 MHz, $\text{DMSO}-d_6$): δ 161.5, 157.7, 152.5, 152.4, 150.2, 135.9, 133.5, 132.9, 131.9, 130.6, 128.9, 126.4, 121.7, 114.2, 55.8. Anal. Calcd for $\text{C}_{17}\text{H}_{12}\text{BrN}_5\text{O}_2$ (398.21): C, 51.27; H, 3.04; N, 17.59. Found: C, 51.50; H, 3.21; N, 17.58.

6-(2-Bromophenyl)-N-(3,4-dimethoxyphenyl)-[1,2,5]-oxadiazolo[3,4-*b*]pyrazin-5-amine (4e). Dark red powder; yield 109 mg (51%); mp 166–167 °C. R_f 0.20 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ 9.29 (s, 1H), 7.97–7.84 (m, 1H), 7.75 (dd, $J = 7.6$, 1.8 Hz, 1H), 7.66 (td, $J = 7.6$, 1.1 Hz, 1H), 7.59 (td, $J = 7.8$, 1.8 Hz, 1H), 7.29–7.22 (m, 1H), 7.15 (d, $J = 2.4$ Hz, 1H), 7.01 (d, $J = 8.7$ Hz, 1H), 3.78 (s, 3H), 3.73 (s, 3H). ^{13}C NMR (151 MHz, $\text{DMSO}-d_6$): δ 161.4, 152.5, 152.3, 150.2, 148.7, 147.4, 135.8, 133.5, 132.9, 131.9, 130.8, 128.9, 121.7, 117.2, 111.9, 109.3, 56.13, 56.11. Anal. Calcd for $\text{C}_{18}\text{H}_{14}\text{BrN}_5\text{O}_3$ (428.24): C, 50.48; H, 3.30; N, 16.35. Found: C, 50.57; H, 3.28; N, 16.07.

6-(2-Bromophenyl)-N-(3-nitrophenyl)-[1,2,5]oxadiazolo[3,4-*b*]pyrazin-5-amine (4g). Dark orange powder; yield 52

mg (25%); mp 163–165 °C. R_f 0.15 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ 9.75 (s, 1H), 8.58 (t, $J = 2.2$ Hz, 1H), 8.23 (ddd, $J = 8.2$, 2.1, 0.9 Hz, 1H), 8.10 (ddd, $J = 8.2$, 2.3, 0.9 Hz, 1H), 7.88 (dd, $J = 8.1$, 1.1 Hz, 1H), 7.78 (dd, $J = 7.6$, 1.7 Hz, 1H), 7.74 (t, $J = 8.2$ Hz, 1H), 7.69 (td, $J = 7.5$, 1.1 Hz, 1H), 7.63 (td, $J = 7.8$, 1.8 Hz, 1H). ^{13}C NMR (151 MHz, $\text{DMSO}-d_6$): δ 161.0, 152.2, 151.9, 150.3, 148.2, 139.2, 135.6, 133.6, 133.1, 132.2, 130.4, 130.4, 128.9, 121.8, 120.6, 118.7. Anal. Calcd for $\text{C}_{16}\text{H}_9\text{BrN}_5\text{O}_3$ (413.19): C, 46.51; H, 2.20; N, 20.34. Found: C, 46.36; H, 2.19; N, 20.07.

6-(2-Bromophenyl)-N-(4-nitrophenyl)-[1,2,5]oxadiazolo[3,4-*b*]pyrazin-5-amine (4h). Orange powder; yield 79 mg (38%); mp 200–202 °C. R_f 0.23 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ 9.81 (s, 1H), 8.36–8.29 (m, 2H), 8.06–8.01 (m, 2H), 7.87 (dd, $J = 8.1$, 1.1 Hz, 1H), 7.78 (dd, $J = 7.6$, 1.7 Hz, 1H), 7.69 (td, $J = 7.5$, 1.1 Hz, 1H), 7.62 (td, $J = 7.8$, 1.8 Hz, 1H). ^{13}C NMR (151 MHz, $\text{DMSO}-d_6$): δ 161.0, 151.9, 151.8, 150.3, 144.3, 144.2, 135.6, 133.6, 133.1, 132.4, 128.9, 124.8, 123.9, 121.8. Anal. Calcd for $\text{C}_{16}\text{H}_9\text{BrN}_5\text{O}_3$ (413.19): C, 46.51; H, 2.20; N, 20.34. Found: C, 46.71; H, 2.09; N, 20.36.

6-(2-Bromophenyl)-N-(*m*-tolyl)-[1,2,5]oxadiazolo[3,4-*b*]pyrazin-5-amine (4i). Yellow powder; yield 102 mg (51%); mp 169–170 °C. R_f 0.50 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (500 MHz, $\text{DMSO}-d_6$): δ 9.35 (s, 1H), 7.85 (dd, $J = 8.1$, 1.1 Hz, 1H), 7.75 (dd, $J = 7.6$, 1.8 Hz, 1H), 7.66 (td, $J = 7.5$, 1.1 Hz, 1H), 7.58 (td, $J = 7.8$, 1.8 Hz, 1H), 7.50–7.44 (m, 1H), 7.37 (d, $J = 2.0$ Hz, 1H), 7.31 (t, $J = 7.8$ Hz, 1H), 7.07 (ddt, $J = 7.5$, 1.8, 0.9 Hz, 1H), 2.32 (s, 3H). ^{13}C NMR (126 MHz, $\text{DMSO}-d_6$): δ 160.9, 151.9, 151.8, 149.7, 137.8, 137.2, 135.3, 133.0, 132.4, 131.5, 128.4, 128.3, 126.6, 124.7, 121.4, 121.2, 21.0. Anal. Calcd for $\text{C}_{17}\text{H}_{12}\text{BrN}_5\text{O}_2$ (398.21): C, 51.27; H, 3.04; N, 17.59. Found: C, 51.37; H, 3.25; N, 17.54.

General Procedure for the Buchwald–Hartwig Cross-Coupling Reaction for the Synthesis of Compounds 6a and 7a.

A stirred mixture of 5-(3-bromothiophen-2-yl)-[1,2,5]oxadiazolo[3,4-*b*]pyrazine (5a) (142 mg, 0.5 mmol), aniline (2a) (55 mg, 0.6 mmol), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (Xantphos) (58 mg, 20 mol %), $\text{Pd}(\text{OAc})_2$ (11 mg, 10 mol %), and K_3PO_4 (265 mg, 1.25 mmol) in degassed 1,4-dioxane (15 mL) was heated at reflux under nitrogen for 15 h in a Schlenk tube. The reaction mixture was cooled, filtered, and dissolved with a mixture of AcOEt and water 1:1 (50 mL), and the organic layer was separated. The aqueous layer was extracted with AcOEt (2 \times 25 mL). The combined organic extracts were dried with MgSO_4 and the solvents were evaporated. The organic layer was separated, dried over anhydrous Na_2SO_4 , and concentrated. The crude products 6a and 7a were purified by column chromatography using hexane and ethyl acetate as an eluent.

2-([1,2,5]Oxadiazolo[3,4-*b*]pyrazin-5-yl)-N-phenylthiophen-3-amine (6a). Violet powder; yield 15 mg (10%); mp 167–169 °C. R_f 0.12 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v). ^1H NMR (500 MHz, $\text{DMSO}-d_6$): δ 10.83 (s, 1H), 9.19 (s, 1H), 8.18 (d, $J = 5.5$ Hz, 1H), 7.46–7.42 (m, 2H), 7.36–7.34 (m, 2H), 7.32 (d, $J = 5.5$ Hz, 1H), 7.18 (tt, $J = 7.2$, 1.2 Hz, 1H). ^{13}C NMR (126 MHz, $\text{DMSO}-d_6$): δ 157.2, 154.2, 152.1, 151.3, 150.7, 140.1, 137.4, 129.7, 124.1, 120.7, 118.7, 111.7. Anal. Calcd for $\text{C}_{14}\text{H}_9\text{N}_5\text{OS}$ (295.32): C, 56.94; H, 3.07; N, 23.71. Found: C, 56.72; H, 3.01; N, 23.48.

8-Phenyl-8H-[1,2,5]oxadiazolo[3,4-*b*]thieno[2',3':4,5]-pyrrolo[2,3-*e*]pyrazine (7a). Orange powder; yield 109 mg (74%); mp 274–275 °C. R_f 0.21 ($\text{CHCl}_3/\text{hexane} = 1:1$, v/v).

^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ 8.54 (d, J = 5.1 Hz, 1H), 7.80–7.77 (m, 2H), 7.70–7.67 (m, 2H), 7.55 (td, J = 7.3, 1.2 Hz, 1H), 7.42 (d, J = 5.1 Hz, 1H). ^{13}C NMR (151 MHz, $\text{DMSO}-d_6$): δ 159.4, 153.7, 151.6, 151.0, 148.0, 143.9, 135.1, 130.3, 128.7, 125.7, 114.6, 113.0. IR (DRA): 447, 485, 514, 556, 592, 606, 619, 649, 663, 675, 690, 704, 743, 759, 780, 793, 829, 859, 867, 882, 894, 915, 994, 1008, 1031, 1041, 1058, 1078, 1094, 1109, 1162, 1198, 1230, 1244, 1290, 1315, 1360, 1383, 1404, 1449, 1474, 1497, 1545, 1557, 1591, 3093, 3107 cm^{-1} . Anal. Calcd for $\text{C}_{14}\text{H}_7\text{N}_3\text{OS}$ (293.30): C, 57.33; H, 2.41; N, 23.88. Found: C, 57.50; H, 2.47; N, 23.99.

General Procedure for the Buchwald–Hartwig Cross-Coupling Reaction for the Synthesis of Compounds 6b, 6c, and 6d. A stirred mixture of *N*-phenyl-2-(quinoxalin-2-yl)thiophen-3-amine (**5b**) [*N*-phenyl-2-(pyrazin-2-yl)aniline (**5c**) or *N*-phenyl-2-(quinoxalin-2-yl)aniline (**5d**) (0.5 mmol), aniline (**2a**) (55 mg, 0.6 mmol), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (Xantphos) (58 mg, 20 mol %), $\text{Pd}(\text{OAc})_2$ (11 mg, 10 mol %), and K_3PO_4 (265 mg, 1.25 mmol) in degassed 1,4-dioxane (15 mL) was heated at reflux under nitrogen for 15 h in a Schlenk tube. The reaction mixture was cooled, filtered, and dissolved with a mixture of AcOEt and water 1:1 (50 mL), and the organic layer was separated. The aqueous layer was extracted with AcOEt (2 \times 25 mL). The combined organic extracts were dried with MgSO_4 and the solvents were evaporated. The organic layer was separated, dried over anhydrous Na_2SO_4 , and concentrated. The crude products **6b–d** were purified by column chromatography using hexane and ethyl acetate as an eluent.

***N*-Phenyl-2-(quinoxalin-2-yl)thiophen-3-amine (6b).** Orange powder; yield 135 mg (89%); mp 148–150 $^\circ\text{C}$. R_f 0.41 ($\text{CHCl}_3/\text{hexane}$ = 1:1, v/v). ^1H NMR (500 MHz, $\text{DMSO}-d_6$): δ 10.38 (s, 1H), 9.11 (s, 1H), 8.10 (dd, J = 8.3, 1.3 Hz, 1H), 8.01 (dd, J = 8.2, 1.4 Hz, 1H), 7.85 (d, J = 5.4 Hz, 1H), 7.82 (ddd, J = 8.3, 6.9, 1.5 Hz, 1H), 7.72 (ddd, J = 8.3, 6.9, 1.4 Hz, 1H), 7.37–7.32 (m, 3H), 7.27–7.24 (m, 2H), 6.99 (tt, J = 7.3, 1.2 Hz, 1H). ^{13}C NMR (126 MHz, $\text{DMSO}-d_6$): δ 148.8, 145.5, 143.9, 142.4, 140.2, 139.3, 130.7, 130.1, 129.5, 128.7, 128.4, 127.8, 121.4, 120.8, 117.9, 113.5. Anal. Calcd for $\text{C}_{18}\text{H}_{13}\text{N}_3\text{S}$ (303.38): C, 71.26; H, 4.32; N, 13.85. Found: C, 71.07; H, 4.23; N, 13.84.

***N*-Phenyl-2-(pyrazin-2-yl)aniline (6c).** Yellow powder; yield 98 mg (79%); mp 82–84 $^\circ\text{C}$. R_f 0.37 ($\text{CHCl}_3/\text{hexane}$ = 1:1, v/v). ^1H NMR (500 MHz, $\text{DMSO}-d_6$): δ 9.22 (s, 1H), 9.07 (d, J = 1.5 Hz, 1H), 8.72 (t, J = 2.0 Hz, 1H), 8.57 (d, J = 2.6 Hz, 1H), 7.78 (d, J = 7.7 Hz, 1H), 7.38 (d, J = 4.0 Hz, 2H), 7.22 (t, J = 7.8 Hz, 2H), 7.10–7.02 (m, 3H), 6.86 (t, J = 7.3 Hz, 1H). ^{13}C NMR (126 MHz, $\text{DMSO}-d_6$): δ 153.1, 144.4, 142.9, 142.8, 142.1, 142.0, 130.5, 130.5, 129.2, 124.3, 120.8, 120.6, 118.5, 117.9. Anal. Calcd for $\text{C}_{16}\text{H}_{13}\text{N}_3$ (247.29): C, 77.71; H, 5.30; N, 16.99. Found: C, 77.72; H, 5.29; N, 17.02.

***N*-Phenyl-2-(quinoxalin-2-yl)aniline (6d).** Yellow powder; yield 144 mg (77%); mp 122–123 $^\circ\text{C}$. R_f 0.23 ($\text{CHCl}_3/\text{hexane}$ = 1:1, v/v). ^1H NMR (500 MHz, $\text{DMSO}-d_6$): δ 9.93 (s, 1H), 9.38 (s, 1H), 8.22 (dd, J = 8.2, 1.5 Hz, 1H), 8.09 (dd, J = 8.2, 1.5 Hz, 1H), 8.03 (dd, J = 7.9, 1.5 Hz, 1H), 7.87 (dddd, J = 22.6, 8.4, 7.0, 1.5 Hz, 2H), 7.49–7.42 (m, 2H), 7.27 (dd, J = 8.5, 7.2 Hz, 2H), 7.20–7.15 (m, 2H), 7.10 (ddd, J = 8.1, 7.0, 1.4 Hz, 1H), 6.90 (tt, J = 7.2, 1.2 Hz, 1H). ^{13}C NMR (126 MHz, $\text{DMSO}-d_6$): δ 152.7, 145.8, 142.7, 142.5, 140.2, 139.9, 131.1, 130.9, 130.5, 129.7, 129.3, 128.7, 128.6, 123.6, 120.9, 120.5, 118.1, 117.9. Anal. Calcd for $\text{C}_{20}\text{H}_{15}\text{N}_3$ (297.35): C, 80.78; H, 5.08; N, 14.13. Found: C, 80.70; H, 5.02; N, 14.20.

Synthesis of 4-Phenyl-4*H*-thieno[2',3':4,5]pyrrolo-[2,3-*b*]quinoxaline (7b). The suspension of *N*-phenyl-2-(quinoxalin-2-yl)thiophen-3-amine (**6b**) (152 mg, 0.5 mmol) in trifluoroacetic acid (5 mL) was stirred at 50 $^\circ\text{C}$ for 24 h. The solvent was removed under reduced pressure. The crude product **7b** was purified by column chromatography using hexane and ethyl acetate as an eluent.

4-Phenyl-4*H*-thieno[2',3':4,5]pyrrolo[2,3-*b*]quinoxaline (7b). Dark orange powder; yield 122 mg (81%); mp 200–201 $^\circ\text{C}$. R_f 0.61 ($\text{CHCl}_3/\text{hexane}$ = 1:1, v/v). ^1H NMR (500 MHz, CDCl_3): δ 8.03 (d, J = 8.0 Hz, 1H), 7.85 (dd, J = 40.7, 8.1 Hz, 2H), 7.52 (dt, J = 21.7, 7.3 Hz, 2H), 7.31–7.21 (m, 2H), 7.07 (t, J = 7.5 Hz, 1H), 6.55 (dd, J = 62.2, 6.7 Hz, 2H), 5.96 (s, 1H). ^{13}C NMR (126 MHz, CDCl_3): δ 150.1, 147.6, 147.2, 141.6, 141.4, 133.5, 131.2, 130.4, 129.7, 129.0, 128.4, 127.9, 126.7, 123.9, 121.7, 119.1. IR (DRA): 511, 538, 572, 607, 647, 698, 718, 737, 753, 789, 804, 841, 866, 894, 944, 974, 1017, 1048, 1062, 1100, 1128, 1139, 1164, 1190, 1209, 1234, 1272, 1313, 1348, 1380, 1402, 1430, 1446, 1482, 1516, 1529, 1577, 1596, 1618, 3057, 3099, 3181, 3279 cm^{-1} . Anal. Calcd for $\text{C}_{18}\text{H}_{11}\text{N}_3\text{S}$ (301.37): C, 71.74; H, 3.68; N, 13.94. Found: C, 71.86; H, 3.35; N, 14.07.

General Procedure for the Synthesis of 5-Phenyl-5*H*-pyrazino[2,3-*b*]indole (7c) and 6-Phenyl-6*H*-indolo[2,3-*b*]quinoxaline (7d). The suspension of *N*-phenyl-2-(pyrazin-2-yl)aniline (**6c**) [or *N*-phenyl-2-(quinoxalin-2-yl)aniline (**6d**)] (0.5 mmol) in EtOH/conc. HCl (10 mL, v/v 99:1) was refluxed for 1 h. The solvent was removed under reduced pressure. The crude products **7c** and **7d** were purified by column chromatography using hexane and ethyl acetate as an eluent.

5-Phenyl-5*H*-pyrazino[2,3-*b*]indole (7c). Beige powder; yield 98 mg (80%); mp 158–160 $^\circ\text{C}$. R_f 0.52 ($\text{EtOAc}/\text{hexane}$ = 1:2, v/v). ^1H NMR (500 MHz, CDCl_3): δ 8.56 (d, J = 2.7 Hz, 1H), 8.43 (dt, J = 7.8, 1.0 Hz, 1H), 8.40 (d, J = 2.7 Hz, 1H), 7.65–7.60 (m, 4H), 7.59 (dd, J = 7.0, 1.3 Hz, 1H), 7.55 (d, J = 8.1 Hz, 1H), 7.50 (tt, J = 5.8, 2.6 Hz, 1H), 7.44 (ddd, J = 8.0, 7.0, 1.2 Hz, 1H). ^{13}C NMR (126 MHz, CDCl_3): δ 145.8, 141.4, 139.8, 137.6, 136.6, 135.3, 129.8, 129.3, 128.1, 127.1, 121.8, 121.7, 119.9, 110.7. IR (DRA): 449, 492, 533, 569, 594, 619, 638, 655, 695, 715, 740, 748, 762, 794, 804, 845, 866, 906, 936, 951, 973, 1012, 1024, 1077, 1104, 1149, 1157, 1171, 1211, 1230, 1270, 1281, 1304, 1327, 1344, 1407, 1456, 1477, 1503, 1545, 1579, 1596, 1621, 3055, 3072 cm^{-1} . Anal. Calcd for $\text{C}_{16}\text{H}_{11}\text{N}_3$ (245.28): C, 78.35; H, 4.52; N, 17.13. Found: C, 78.34; H, 4.76; N, 17.16.

6-Phenyl-6*H*-indolo[2,3-*b*]quinoxaline (7d). Pale yellow powder; yield 109 mg (74%); mp 237–238 $^\circ\text{C}$. R_f 0.32 ($\text{CHCl}_3/\text{hexane}$ = 1:1, v/v). ^1H NMR (500 MHz, CDCl_3): δ 8.57–8.51 (m, 1H), 8.32 (dd, J = 7.9, 1.9 Hz, 1H), 8.09 (dd, J = 7.9, 1.9 Hz, 1H), 7.77–7.62 (m, 7H), 7.54–7.49 (m, 2H), 7.44 (t, J = 7.5 Hz, 1H). ^{13}C NMR (126 MHz, CDCl_3): δ 145.8, 144.7, 140.5, 140.1, 139.8, 135.4, 131.0, 129.8, 129.2, 128.8, 128.2, 127.9, 127.1, 126.5, 122.6, 121.8, 119.8, 110.6. IR (DRA): 591, 619, 650, 695, 720, 749, 767, 781, 802, 830, 859, 909, 924, 955, 1007, 1016, 1027, 1043, 1074, 1101, 1134, 1148, 1175, 1207, 1228, 1254, 1305, 1320, 1337, 1355, 1392, 1405, 1460, 1471, 1484, 1504, 1583, 1598, 1609, 1629, 3057 cm^{-1} . Anal. Calcd for $\text{C}_{20}\text{H}_{13}\text{N}_3$ (295.34): C, 81.34; H, 4.44; N, 14.23. Found: C, 81.30; H, 4.51; N, 14.21.

■ ASSOCIATED CONTENT

■ Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acsomega.0c01945>.

Crystallographic data for **3i** (CIF)

Experimental procedures, UV–vis absorption spectra of compounds **3a**, **3d**, **3h**, and **7a–d**, characterization data for all new compounds, and ^1H and ^{13}C NMR spectra (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Egor V. Verbitskiy – I. Postovsky Institute of Organic Synthesis, Ural Branch of the Russian Academy of Sciences, Ekaterinburg 620108, Russia; Ural Federal University, Ekaterinburg 620002, Russia; orcid.org/0000-0002-9613-8738; Email: verbitskiy@ios.uran.ru

Authors

Yuriy A. Kvashnin – I. Postovsky Institute of Organic Synthesis, Ural Branch of the Russian Academy of Sciences, Ekaterinburg 620108, Russia

Ekaterina F. Zhilina – I. Postovsky Institute of Organic Synthesis, Ural Branch of the Russian Academy of Sciences, Ekaterinburg 620108, Russia

Gennady L. Rusinov – I. Postovsky Institute of Organic Synthesis, Ural Branch of the Russian Academy of Sciences, Ekaterinburg 620108, Russia; Ural Federal University, Ekaterinburg 620002, Russia

Oleg N. Chupakhin – I. Postovsky Institute of Organic Synthesis, Ural Branch of the Russian Academy of Sciences, Ekaterinburg 620108, Russia; Ural Federal University, Ekaterinburg 620002, Russia; orcid.org/0000-0002-1672-2476

Valery N. Charushin – I. Postovsky Institute of Organic Synthesis, Ural Branch of the Russian Academy of Sciences, Ekaterinburg 620108, Russia; Ural Federal University, Ekaterinburg 620002, Russia

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/acsomega.0c01945>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by the Russian Foundation for Basic Research (research project no. 18-29-23045 mk). Y.A.K. would like to acknowledge the financial support for the part of the synthetic section from the Russian Foundation for Basic Research (research project no. 18-33-00103-mol_a).

■ REFERENCES

- (1) (a) Pai, N. R.; Pusalkar, D. A. Pharmacological screening of novel indolo[2,3-*b*]quinoxaline derivatives. *J. Chem. Pharm. Res.* **2010**, *2*, 485–493. (b) Manna, K.; Agrawal, Y. K. Microwave assisted synthesis of new indophenazine 1,3,5-trisubstituted pyrazoline derivatives of benzofuran and their antimicrobial activity. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 2688–2692.
- (2) (a) Hirata, K.; Araya, J.; Nakaike, S.; Kitamura, K.; Ishida, T. Side Chain-Dependent Binding of Antitumor Indoloquinoxaline Derivatives to DNA: Comparative Spectroscopic and Viscometric Measurements. *Chem. Pharm. Bull.* **2001**, *49*, 44–48. (b) Deady, L. W.; Kaye, A. J.; Finlay, G. J.; Baguley, B. C.; Denny, W. A. Synthesis and Antitumor Properties of *N*-[2-(Dimethylamino)ethyl]-carboxamide Derivatives of Fused Tetracyclic Quinolines and Quinoxalines: A New Class of Putative Topoisomerase Inhibitors. *J. Med. Chem.* **1997**, *40*, 2040–2046. (c) Avula, S.; Komsani, J. R.; Koppireddi, S.; Yadla, R.; Kanugula, A. K.; Kotamraju, S. Synthesis and cytotoxicity of novel 6*H*-indolo[2,3-*b*]quinoxaline derivatives. *Med. Chem. Res.* **2013**, *22*, 3712–3718.
- (3) (a) Harmenberg, J.; Wahren, B.; Bergman, J.; Akerfeldt, S.; Lundblad, L. Antiherpesvirus activity and mechanism of action of indolo-(2,3-*b*)quinoxaline and analogs. *Antimicrob. Agents Chemother.* **1988**, *32*, 1720–1724. (b) Harmenberg, J.; Åkesson-Johansson, A.; Gräslund, A.; Malmfors, T.; Bergman, J.; Wahren, B.; Åkerfeldt, S.; Lundblad, L.; Cox, S. The mechanism of action of the anti-herpes virus compound 2,3-dimethyl-6(2-dimethylaminoethyl)-6*H*-indolo-(2,3-*b*)quinoxaline. *Antiviral Res.* **1991**, *15*, 193–204.
- (4) (a) Su, T.-H.; Fan, C.-H.; Ou-Yang, Y.-H.; Hsu, L.-C.; Cheng, C.-H. Highly efficient deep-red organic electrophosphorescent devices with excellent operational stability using bis(indoloquinoxaliny) derivatives as the host materials. *J. Mater. Chem. C* **2013**, *1*, 5084–5092. (b) Tyagi, P.; Venkateswararao, A.; Thomas, K. R. J. Solution Processable Indoloquinoxaline Derivatives Containing Bulky Polyaromatic Hydrocarbons: Synthesis, Optical Spectra, and Electroluminescence. *J. Org. Chem.* **2011**, *76*, 4571–4581. (c) Fan, C.-H.; Sun, P.; Su, T.-H.; Cheng, C.-H. Host and Dopant Materials for Idealized Deep-Red Organic Electrophosphorescence Devices. *Adv. Mater.* **2011**, *23*, 2981–2985.
- (5) Thomas, K. R. J.; Tyagi, P. Synthesis, Spectra, and Theoretical Investigations of the Triarylamines Based on 6*H*-Indolo[2,3-*b*]quinoxaline. *J. Org. Chem.* **2010**, *75*, 8100–8111.
- (6) Zhao, J.; Li, H.; Li, H.; Zhao, Q.; Ling, H.; Li, J.; Lin, J.; Xie, L.; Lin, Z.; Yi, M.; Huang, W. Synthesis, characterization and charge storage properties of π -biindolo[2,3-*b*]quinoxaline for solution-processing organic transistor memory. *Dyes Pigm.* **2019**, *167*, 255–261.
- (7) (a) Dandia, A.; Parewa, V.; Maheshwari, S.; Rathore, K. S. Cu doped CdS nanoparticles: A versatile and recoverable catalyst for chemoselective synthesis of indolo[2,3-*b*]quinoxaline derivatives under microwave irradiation. *J. Mol. Catal. A: Chem.* **2014**, *394*, 244–252. (b) Edayadulla, N.; Lee, Y. R. Cerium oxide nanoparticle-catalyzed three-component protocol for the synthesis of highly substituted novel quinoxalin-2-amine derivatives and 3,4-dihydroquinoxalin-2-amines in water. *RSC Adv.* **2014**, *4*, 11459–11468.
- (8) Mannes, P. Z.; Onyango, E. O.; Gribble, G. W. Synthesis of a Masked 2,3-Diaminoindole. *J. Org. Chem.* **2016**, *81*, 12478–12481.
- (9) (a) Ackermann, L.; Althammer, A.; Mayer, P. Palladium-Catalyzed Direct Arylation-Based Domino Synthesis of Annulated *N*-Heterocycles Using Alkenyl or (Hetero)Aryl 1,2-Dihalides. *Synthesis* **2009**, *2009*, 3493–3503. (b) Ackermann, L.; Althammer, A. Domino *N*-H/C-H bond activation: palladium-catalyzed synthesis of annulated heterocycles using dichloro(hetero)arenes. *Angew. Chem., Int. Ed.* **2007**, *46*, 1627–1629. (c) Hung, T. Q.; Hoang, D. H.; Thang, N. N.; Dang, T. T.; Ayub, K.; Villinger, A.; Friedrich, A.; Lochbrunner, S.; Flechsig, G.-U.; Langer, P. Palladium catalyzed synthesis and physical properties of indolo[2,3-*b*]quinoxalines. *Org. Biomol. Chem.* **2014**, *12*, 6151–6166.
- (10) Hung, T. Q.; Dang, T. T.; Villinger, A.; Sung, T. V.; Langer, P. Efficient synthesis of thieno[3,2-*b*:4,5-*b'*]diindoles and benzothieno[3,2-*b*]indoles by Pd-catalyzed site-selective C–C and C–N coupling reactions. *Org. Biomol. Chem.* **2012**, *10*, 9041–9044.
- (11) (a) Trost, B. The atom economy—a search for synthetic efficiency. *Science* **1991**, *254*, 1471–1477. (b) Trost, B. M. Atom Economy—A Challenge for Organic Synthesis: Homogeneous Catalysis Leads the Way. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 259–281.
- (12) Baek, Y.; Maeng, C.; Kim, H.; Lee, P. H. Regioselective Synthesis of Indolopyrazines through a Sequential Rhodium-Catalyzed Formal [3+3] Cycloaddition and Aromatization Reaction of Diazoindolinimines with Azirines. *J. Org. Chem.* **2018**, *83*, 2349–2360.

(13) For selected reviews on the metal-free oxidative C–H bond functionalization, see: (a) Chupakhin, O. N.; Charushin, V. N.; Van der Plas, H. C. *Nucleophilic Aromatic Substitution of Hydrogen*; Academic Press: New York, San Diego, 1994. (b) Charushin, V. N.; Chupakhin, O. N. S_N^H methodology and new approaches to condensed heterocyclic systems. *Pure Appl. Chem.* **2004**, *76*, 1621–1631. (c) Charushin, V. N.; Chupakhin, O. N. Nucleophilic aromatic substitution of hydrogen and related reactions. *Mendeleev Commun.* **2007**, *17*, 249–254. (d) Makosza, M.; Wojciechowski, K. Nucleophilic Substitution of Hydrogen in Heterocyclic Chemistry. *Chem. Rev.* **2004**, *104*, 2631–2666. (e) Makosza, M. Nucleophilic substitution of hydrogen in electron-deficient arenes, a general process of great practical value. *Chem. Soc. Rev.* **2010**, *39*, 2855–2868. (f) *Topics in Heterocyclic Chemistry*; Charushin, V. N., Chupakhin, O. N., Eds.; Springer International Publishing: Switzerland, 2013; pp 1–50. (g) Makosza, M.; Wojciechowski, K. In *Topics in Heterocyclic Chemistry*; Charushin, V. N., Chupakhin, O. N., Eds.; Springer International Publishing: Switzerland, 2013; pp 51–105. (h) Verbitskiy, E.; Rusinov, G.; Chupakhin, O.; Charushin, V. Recent Advances in Direct C–H Functionalization of Pyrimidines. *Synthesis* **2018**, *50*, 193–210.

(14) (a) Sheremetev, A. B.; Yudin, I. L. Advances in the chemistry of furazano[3,4-*b*]pyrazines and their analogues. *Russ. Chem. Rev.* **2003**, *72*, 87–100. (b) Kvashnin, Y. A.; Kazin, N. A.; Verbitskiy, E. V.; Svalova, T. S.; Ivanova, A. V.; Kozitsina, A. N.; Slepukhin, P. A.; Rusinov, G. L.; Chupakhin, O. N.; Charushin, V. N. A facile, metal-free, oxidative coupling of new 6-(hetero)aryl-[1,2,5]oxadiazolo[3,4-*b*]pyrazines with pyrroles, indoles and carbazoles. *ARKIVOC* **2016**, *2016*, 279–300. (c) Verbitskiy, E. V.; Kvashnin, Y. A.; Slepukhin, P. A.; Kuchin, A. V.; Rusinov, G. L.; Chupakhin, O. N.; Charushina, V. N. Reactions of pyrazinium salts with phenols: from σ^H -adducts to S_N^H products and transformations into benzo[*b*]furans. *Russ. Chem. Bull.* **2011**, *60*, 919–928. (d) Verbitskiy, E. V.; Kvashnin, Y. A.; Baranova, A. A.; Yakovleva, Y. A.; Khokhlov, K. O.; Rusinov, G. L.; Charushin, V. N. 9-Ethyl-3-{6-(het)aryl-[1,2,5]oxadiazolo[3,4-*b*]pyrazin-5-yl}-9H-carbazoles: synthesis and study of sensitivity to nitroaromatic compounds. *Russ. Chem. Bull.* **2018**, *67*, 1078–1082. (e) Utepova, I. A.; Musikhina, A. A.; Kvashnin, Y. A.; Shcherbakova, M. A.; Serebrennikova, P. O.; Rusinov, G. L.; Chupakhin, O. N. Direct C–C-coupling of ferrocenyl- and cymantrenyllithium with 5-(het)aryl-1,2,5-oxadiazolo[3,4-*b*]pyrazines. *Russ. Chem. Bull.* **2011**, *60*, 2531–2535. (f) Kazin, N. A.; Kvashnin, Y. A.; Irgashev, R. A.; Dehaen, W.; Rusinov, G. L.; Charushin, V. N. Direct arylalkenylation of furazano[3,4-*b*]pyrazines via a new C–H functionalization protocol. *Tetrahedron Lett.* **2015**, *56*, 1865–1869. (g) Tsyganov, D. V.; Samet, A. V.; Dorovatovskii, P. V.; Khrustalev, V. N.; Semenov, V. V. Arylglyoxal oximes as putative C-nucleophiles in eliminative nucleophilic substitution process. *Mendeleev Commun.* **2019**, *29*, 296–298.

(15) Catrow, J. L.; Zhang, Y.; Zhang, M.; Ji, H. Discovery of Selective Small-Molecule Inhibitors for the β -Catenin/T-Cell Factor Protein–Protein Interaction through the Optimization of the Acyl Hydrazone Moiety. *J. Med. Chem.* **2015**, *58*, 4678–4692.

(16) Verbitskiy, E. V.; Cheprakova, E. M.; Slepukhin, P. A.; Kodess, M. I.; Ezhikova, M. A.; Pervova, M. G.; Rusinov, G. L.; Chupakhin, O. N.; Charushin, V. N. Combination of the Suzuki–Miyaura cross-coupling and nucleophilic aromatic substitution of hydrogen (S_N^H) reactions as a versatile route to pyrimidines bearing thiophene fragments. *Tetrahedron* **2012**, *68*, 5445–5452.

(17) Ponizovskiy, M. G.; Boguslavsky, A. M.; Kodess, M. I.; Charushin, V. N.; Chupakhin, O. N. Synthesis of fused quinoxalines. *Mendeleev Commun.* **2002**, *12*, 68–70.

(18) Loukrakpam, D. C.; Phukan, P. $TsNBr_2$ Mediated Synthesis of 2-Acylbenzothiazoles and Quinoxalines from Aryl Methyl Ketones under Metal Free Condition. *ChemistrySelect* **2019**, *4*, 3180–3184.